

Perturbing the hexagonal circle packing: a percolation perspective

Itai Benjamini*

Alexandre Stauffer†

January 19, 2013

Abstract

We consider the hexagonal circle packing with radius $1/2$ and perturb it by letting the circles move as independent Brownian motions for time t . It is shown that, for large enough t , if Π_t is the point process given by the center of the circles at time t , then, as $t \rightarrow \infty$, the critical radius for circles centered at Π_t to contain an infinite component converges to that of continuum percolation (which was shown—based on a Monte Carlo estimate—by Balister, Bollobás and Walters to be strictly bigger than $1/2$). On the other hand, for small enough t , we show (using a Monte Carlo estimate for a fixed but high dimensional integral) that the union of the circles contains an infinite connected component. We discuss some extensions and open problems.

1 Introduction

Let $\mathcal{T}$ be the triangular lattice with edge length 1 and let Π_0 be the set of vertices of $\mathcal{T}$. We see Π_0 as a point process and, to avoid ambiguity, we use the term *node* to refer to the points of Π_0 . Now, for each node $u \in \Pi_0$, we add a ball of radius $1/2$ centered at u , and set $R(\Pi_0)$ to be the region of $\mathbb{R}^2$ obtained by the union of these balls; more formally,

$$R(\Pi_0) = \bigcup_{x \in \Pi_0} B(x, 1/2),$$

where $B(y, r)$ denotes the closed ball of radius r centered at y . In this way, $R(\Pi_0)$ is the so-called hexagonal circle packing of $\mathbb{R}^2$; refer to [4] for more information on packings. Clearly, the region $R(\Pi_0)$ is a *connected* subset of $\mathbb{R}^2$.

Our goal is to analyze how this set evolves as we let the nodes of Π_0 move on $\mathbb{R}^2$ according to independent Brownian motions. For any $t > 0$, let Π_t be the point process obtained after the nodes have moved for time t . More formally, for each node $u \in \Pi_0$, let $(\zeta_u(t))_t$ be an independent Brownian motion on $\mathbb{R}^2$ starting at the origin, and set

$$\Pi_t = \bigcup_{u \in \Pi_0} (u + \zeta_u(t)).$$

A natural question is whether there exists a phase transition on t such that $R(\Pi_t)$ has an infinite component for small t but has only finite components for large t .

Intuitively, for sufficiently large time, one expects that Π_t will look like a Poisson point process with intensity $2/\sqrt{3}$, which is the density of nodes in the triangular lattice. Then, for sufficiently large t , $R(\Pi_t)$ will contain an infinite component almost surely only if $R(\Phi)$ contains an infinite

*Weizmann Institute of Science, Rehovot, Israel Email: itai.benjamini@weizmann.ac.il

†Computer Science Division, University of California, Berkeley CA, USA Email: stauffer@cs.berkeley.edu. Supported by a Fulbright/CAPES scholarship and NSF grants CCF-0635153 and DMS-0528488.

component where Φ is a Poisson point process with intensity $\lambda = 2/\sqrt{3}$. In the literature, $R(\Phi)$ is referred to as the Boolean model. For this model, it is known that there exists a value λ_c so that, if $\lambda < \lambda_c$, then all connected components of $R(\Phi)$ are finite almost surely [9]. On the other hand, if $\lambda > \lambda_c$, then $R(\Phi)$ contains an infinite connected component. The value of λ_c is currently unknown and depends on the radius of the balls in the definition of the region R . When the balls have radius $1/2$, it is known that λ_c satisfies $0.52 \leq \lambda_c \leq 3.38$ [3, Chapter 8], but these bounds do not answer whether $2/\sqrt{3}$ is smaller or larger than λ_c . However, using a Monte Carlo analysis, Balister, Bollobás and Walters [2] showed that, with 99.99% confidence, λ_c lies between 1.434 and 1.438, which are both larger than $2/\sqrt{3}$. We then have the following theorem, whose proof we give in Section 3.

Theorem 1.1. *If $\lambda_c > 2/\sqrt{3}$, where λ_c is the critical intensity for percolation of the Boolean model with balls of radius $1/2$, then there exists a positive t_0 so that, for all $t > t_0$, $R(\Pi_t)$ contains no infinite component almost surely.*

We now give a different statement of Theorem 1.1. For each t , there exists a critical radius $r_c(t)$ so that, adding balls of radius $r > r_c(t)$ centered at the points of Π_t gives that the union of these balls contains an infinite component almost surely. Then, the next theorem, whose proof we give in Section 3, follows from the proof of Theorem 1.1 and a result of Sinclair and Stauffer [13].

Theorem 1.2. *As $t \rightarrow \infty$, we have that $r_c(t)$ converges to the critical radius for percolation of the Boolean model with intensity $2/\sqrt{3}$.*

In order to prove Theorems 1.1 and 1.2, we devote most of Section 3 to prove Theorem 1.3 below, which we believe to be of independent interest. Consider a tessellation of $\mathbb{R}^2$ into regular hexagons of side length $\delta\sqrt{t}$, where δ is an arbitrarily small constant. Let I denote the set of points of $\mathbb{R}^2$ that are the centers of these hexagons. Then, for each $i \in I$, denote the hexagon with center at i by Q_i and define a Bernoulli random variable X_i with parameter p independently of the other X_j , $j \neq i$. Define

$$C(p, \delta) = \bigcup_{i \in I: X_i=1} Q_i.$$

When $p > 1/2$, which is the critical probability for site percolation on the triangular lattice, we have that $C(p, \delta)$ contains a unique infinite connected component. We are now ready to state our main technical result, whose proof is given in Section 3.

Theorem 1.3. *There exists a universal constant $c > 0$ and, for any $p \in (0, 1)$ that can be arbitrarily close to 1 and any arbitrarily small $\delta > 0$, there exists a $t_0 > 0$ such that, for all $t > t_0$, we can couple Π_t , $(X_i)_{i \in I}$ and a Poisson point process Φ of intensity $\frac{2}{\sqrt{3}} + c\sqrt{\delta}$, which is independent of Π_0 , so that*

$$R(\Pi_t) \cap C(p, \delta) \subset R(\Phi) \cap C(p, \delta).$$

In words, Theorem 1.3 establishes that Π_t is *stochastically dominated* by a Poisson point process inside $C(p, \delta)$. As we show later in Lemma 2.1, Π_t cannot be stochastically dominated by a Poisson point process in the whole of $\mathbb{R}^2$.

We note that the opposite direction of Theorem 1.3 was established by Sinclair and Stauffer [13, Proposition 4.1], who proved that, under some conditions on the initial location of the nodes, after moving as independent Brownian motions for time t , the nodes *stochastically dominate* a Poisson point process. The result of Sinclair and Stauffer has been used and refined in [12, 14], and turned out to be a useful tool in the analysis of increasing events for models of mobile nodes, such as the so-called percolation time in [12] and detection time in [14]. We expect that the ideas in our

proof of Theorem 1.3 can help in the analysis of *decreasing* events, which have so far received less attention.

We remark that the proof of our Theorem 1.3 requires a more delicate analysis than that of Sinclair and Stauffer. In their case, nodes that ended up moving atypically far away during the time $[0, t]$ could be simply disregarded as it is possible to show that the typical nodes already stochastically dominate a Poisson point process. In our setting, no node can be disregarded, regardless of how atypical its motion turns out to be. In order to solve this problem, we first consider what we call well-behaved nodes, which among other things satisfy that their motion during $[0, t]$ is contained in some ball of radius $c\sqrt{t}$ for some large constant c (we defer the complete definition of well-behaved nodes to Section 3). The definition of well-behaved nodes is carefully specified so that any given node is likely to be well-behaved and, in addition, it is possible to show that well-behaved nodes are stochastically dominated by a Poisson point process inside $C(p, \delta)$. For the remaining nodes, which comprise only a small density of nodes, we use a sprinkling argument to replace them already at time 0 by a (non-homogeneous) Poisson point process of low intensity. Then, even though the motion of the nodes that are not well behaved is hard to control, we use the fact that they are a Poisson point process at time 0 to show that, at time t , they stochastically dominate a Poisson point process of low intensity.

Now, for the case when the nodes of Π_0 move for only a small time t , we believe the following is true.

Conjecture 1.4. *There exists a $t_0 > 0$ so that, for all $t < t_0$, $R(\Pi_t)$ contains an infinite component almost surely.*

We are able to establish the conjecture above given a Monte Carlo estimate for a finite but high dimensional integral. We discuss the details in Section 4. Note that if Conjecture 1.4 is true, then a curious consequence of this and Theorem 1.2 is that $r_c(t)$ is not monotone in t .

We conclude in Section 5 with some extensions and open problems.

2 Stochastic Domination

We devote this section to the proof of our main technical result, Theorem 1.3, where we study the behavior of the balls after they have moved for a time t that is sufficiently large. In Section 3 we show how to adapt this proof to establish Theorems 1.1 and 1.2.

Intuitively, as $t \rightarrow \infty$, Π_t looks like an independent Poisson point process of intensity $2/\sqrt{3}$. Since the intensity of Φ is larger than $2/\sqrt{3}$, we would like to argue that there exists a coupling between Φ and Π_t such that Φ contains Π_t . Unfortunately, this cannot be achieved in the whole of $\mathbb{R}^2$, as established by the lemma below, which gives that, for any fixed t , the probability that a sufficiently large region $S \subset \mathbb{R}^2$ contains no node of Π_t is smaller than $\exp\left(-\left(\frac{2}{\sqrt{3}} + c\sqrt{\delta}\right) \text{vol}(S)\right)$, which is the probability that Φ has no node in S . Hence, Φ cannot stochastically dominate Π_t in the whole of $\mathbb{R}^2$.

Lemma 2.1. *Fix t sufficiently large and let S be a hexagon of side length $(\log t)\sqrt{t}$ obtained as the union of $(\log t)^2 t$ triangles of $\mathcal{T}$. Then, there exists a positive constant c' such that*

$$\mathbf{P}(\Pi_t \cap S = \emptyset) \leq \exp\left(-c'(\log t)^2 \text{vol}(S)\right).$$

Proof. For simplicity we assume that $(\log t)\sqrt{t}$ is an even integer. Let x be the middle point of S and consider the hexagon S' of side length $\frac{\log t}{2}\sqrt{t}$ composed of $\frac{(\log t)^2}{4}t$ triangles of $\mathcal{T}$ and centered at x . Note that S contains the ball $B\left(x, \frac{\sqrt{3}\log t}{2}\sqrt{t}\right)$ and S' is contained in the ball $B\left(x, \frac{\log t}{2}\sqrt{t}\right)$.

Therefore, a node of Π_0 that is inside S' can only be outside of S at time t if it moves at least $\frac{(\sqrt{3}-1)\log t}{2}\sqrt{t} \geq \frac{\log t}{3}\sqrt{t}$. For any fixed node $u \in \Pi_0$, we have from the Gaussian tail bound (cf. Lemma A.3) that

$$\mathbf{P}\left(\|\zeta_u(t)\|_2 \geq \frac{\log t}{3}\sqrt{t}\right) \leq \frac{3}{\sqrt{2\pi}\log t} \exp\left(-\frac{(\log t)^2}{18}\right).$$

Each node of Π_0 belongs to 6 triangles of $\mathcal{T}$, then there are at least $\frac{(\log t)^2}{24}t$ nodes of Π_0 in S' . Since each of them need to move more than $\frac{\log t}{3}\sqrt{t}$ by time t for S to contain no node of Π_t , we obtain

$$\mathbf{P}(\Pi_t \cap S = \emptyset) \leq \left(\frac{3}{\sqrt{2\pi}\log t} \exp\left(-\frac{(\log t)^2}{18}\right)\right)^{\frac{(\log t)^2}{24}t} \leq \exp\left(-\frac{(\log t)^4 t}{432}\right).$$

Since $\text{vol}(S) = \frac{3\sqrt{3}}{2}(\log t)^2 t$, the proof is completed. $\square$

Now we turn to the proof of Theorem 1.3. The goal is to show that Φ contains Π_t inside a percolating cluster of a suitable tessellation of $\mathbb{R}^2$. For this, we tessellate $\mathbb{R}^2$ into hexagons of side length $\delta\sqrt{t}$. We take this tessellation in such a way that no point of Π_0 lies on the edges of the hexagons; this is not crucial for the proof but simplifies the explanations in the sequel. Let $\mathcal{H}$ denote the set of these hexagons. Consider a node $v \in \Pi_0$. Let Q_i be the hexagon of $\mathcal{H}$ that contains v and let v' be a copy of v located at the same position as v at time 0. We let v' move up to time t according to a certain procedure that we will describe in a moment, and then we say that v is *well behaved* if we are able to couple the motion of v with the motion of v' so that v and v' are at the same location at time t . Recall that I is the set of points given by the centers of the hexagons in $\mathcal{H}$. For $i \in I$, we define

$$J_i = \left\{j \in I: \sup_{x \in Q_i, y \in Q_j} \|x - y\|_2 \leq C\delta\sqrt{t}\right\},$$

where

$$C = 4\delta^{-3/2}.$$

For i, j such that $j \in J_i$ we say that i and j are neighbors. Now we describe the motion of v' . Let f_t be the density function for the location of a Brownian motion at time t given that it starts at the origin of $\mathbb{R}^2$. We fix t and, for each $i, j \in I$ such that i and j are neighbors, we let

$$\varphi_t(i, j) = \inf_{x \in Q_i, y \in Q_j} f_t(y - x). \quad (1)$$

If i and j are not neighbors we set $\varphi_t(i, j) = 0$. Then, the motion of v' is described by first choosing a $j \in J_i$ with probability proportional to $\varphi_t(i, j)$ and then placing v' uniformly at random in Q_j . The main intuition behind this definition is that, when v is well behaved, its position inside Q_j has the same distribution as that of a node of a Poisson point process inside Q_j . Therefore, as long as the number of well behaved nodes that end up in Q_j is smaller than the number of nodes in $\Phi \cap Q_j$, we will be able to couple them with Φ . Another important feature for the definition of well behaved nodes is that, if v is well behaved and ends up moving to hexagon Q_j , then we know that, at time 0, v was in some hexagon of J_j . In particular, there is a bounded number of hexagons from which v could have moved to Q_j , which allows us to control dependences.

Now we show that nodes are likely to be well behaved. Since the area of each hexagon of $\mathcal{H}$ is $\frac{3\sqrt{3}}{2}\delta^2 t$, we have that

$$\mathbf{P}(v \text{ is well behaved}) = \sum_{j \in J_i} \frac{3\sqrt{3}}{2} \delta^2 t \varphi_t(i, j). \quad (2)$$

The idea is that δ is sufficiently small so that f_t varies very little (i.e., f_t is essentially constant) inside any given hexagon of $\mathcal{H}$, but, at the same time, $C\delta$ is large so that the probability that v moves to an hexagon that is not in J_i is small. We can then obtain in the lemma below that the probability that v is well behaved is large.

Lemma 2.2. *Let v be a node of Π_0 located in Q_i . We have*

$$(C - 3)^2 \leq |J_i| \leq \frac{4}{3}C^2,$$

and, for sufficiently large t , we have

$$\mathbf{P}(v \text{ is well behaved}) \geq 1 - 5\delta.$$

Proof. For $j \notin J_i$, we know, by definition, that there exist a $x_0 \in Q_i$ and a $y_0 \in Q_j$ such that $\|x_0 - y_0\|_2 > C\delta\sqrt{t}$. Then, by the triangle inequality, we have that, for any $y \in Q_j$,

$$\|y - i\|_2 \geq C\delta\sqrt{t} - \|y - y_0\|_2 - \|i - x_0\|_2 \geq C\delta\sqrt{t} - 3\delta\sqrt{t}, \quad (3)$$

where we used the fact that, for any two points y, y_0 in the same hexagon, we have $\|y - y_0\|_2 \leq 2\delta\sqrt{t}$ and, for any $x \in Q_i$ we have $\|i - x\|_2 \leq \delta\sqrt{t}$. Therefore, if we add balls of radius $\delta\sqrt{t}$ centered at each $j \in J_i$, these balls cover the whole of $B(i, C\delta\sqrt{t} - 3\delta\sqrt{t})$, which yields

$$|J_i| \geq \frac{\text{vol}(B(i, C\delta\sqrt{t} - 3\delta\sqrt{t}))}{\text{vol}(B(0, \delta\sqrt{t}))} = (C - 3)^2.$$

For the other direction, note that if we add balls of radius $\frac{\sqrt{3}}{2}\delta\sqrt{t}$ centered at each $j \in J_i$, these balls are disjoint and their union is contained in $B(i, C\delta\sqrt{t})$, which gives

$$|J_i| \leq \frac{\text{vol}(B(i, C\delta\sqrt{t}))}{\text{vol}(B(0, \frac{\sqrt{3}}{2}\delta\sqrt{t}))} = \frac{4}{3}C^2.$$

Now we prove the second part of the lemma. Note that, using (2) and (1), we have

$$\begin{aligned} \mathbf{P}(v \text{ is well behaved}) &= \sum_{j \in J_i} \frac{3\sqrt{3}}{2} \frac{\delta^2 t}{2\pi t} \exp\left(-\frac{\sup_{x \in Q_i, y \in Q_j} \|x - y\|_2^2}{2t}\right) \\ &\geq \sum_{j \in J_i} \int_{Q_j} \frac{1}{2\pi t} \exp\left(-\frac{(\|z - i\|_2 + 3\delta\sqrt{t})^2}{2t}\right) dz, \end{aligned} \quad (4)$$

where the last step follows by the triangle inequality. Now, from (3), the ball $B(i, C\delta\sqrt{t} - 3\delta\sqrt{t})$ only intersects hexagons that are neighbors of i . We denote by $S_a = [-a/2, a/2]^2$ the square of side length a , and, for any $z = (z_1, z_2) \in \mathbb{R}^2$ and $a \in \mathbb{R}_+$, we use the inequality $(\|z\|_2 + a)^2 \leq$

$(|z_1| + a)^2 + (|z_2| + a)^2$. Then, applying (4), we obtain

$$\begin{aligned}
\mathbf{P}(v \text{ is well behaved}) &\geq \int_{B(0, C\delta\sqrt{t}-3\delta\sqrt{t})} \frac{1}{2\pi t} \exp\left(-\frac{(\|z\|_2 + 3\delta\sqrt{t})^2}{2t}\right) dz \\
&\geq \int_{S_{\frac{2C\delta\sqrt{t}-6\delta\sqrt{t}}{\sqrt{2}}}} \frac{1}{2\pi t} \exp\left(-\frac{(\|z\|_2 + 3\delta\sqrt{t})^2}{2t}\right) dz \\
&\geq \left(2 \int_{3\delta\sqrt{t}}^{\frac{C\delta\sqrt{t}+3(\sqrt{2}-1)\delta\sqrt{t}}{\sqrt{2}}} \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{z_1^2}{2t}\right) dz_1\right)^2 \\
&\geq \left(1 - \frac{6\delta}{\sqrt{2\pi}} - \frac{2}{\sqrt{\pi}(C+3(\sqrt{2}-1))\delta} \exp\left(-\frac{\delta^2(C+3(\sqrt{2}-1))^2}{4}\right)\right)^2 \\
&\geq 1 - \frac{12\delta}{\sqrt{2\pi}} - \frac{4}{\sqrt{\pi}C\delta} \exp\left(-\frac{\delta^2 C^2}{4}\right),
\end{aligned}$$

where the second to last step follows by the standard Gaussian tail bound (cf. Lemma A.3). Then, using the value of C , we have that $\frac{4}{\sqrt{\pi}C\delta}e^{-\delta^2 C^2/4} \leq \frac{1}{\sqrt{\pi}}e^{-4}$ for all $\delta \in [0, 1]$. Then, the right-hand side above is at least $1 - \delta\left(\frac{12}{\sqrt{2\pi}} + \frac{e^{-4}}{\sqrt{\pi}}\right) \geq 1 - 5\delta$. $\square$

We will treat the nodes that are not well behaved by means of another point process. For any point $x \in \mathbb{R}^2$, we set $q(x) = i$ if $x \in Q_i$. Then, let $g_t(x, y)$ be the density function for a node v that is not well behaved to move from x to y after time t . We have that

$$g_t(x, y) = \frac{f_t(y - x) - \varphi_t(q(x), q(y))}{\mathbf{P}(v \text{ is not well behaved})}. \quad (5)$$

For each $v \in \Pi_0$, let $N_v(\mu)$ be a Poisson random variable with mean μ , and let $\Psi_0(\mu)$ be the point process obtained by putting $N_v(\mu)$ points at v for each $v \in \Pi_0$. We set $e^{-\mu} = \mathbf{P}(v \text{ is well behaved})$ and, from Lemma 2.2 and the fact that δ is sufficiently small, we henceforth assume that $\mu \leq 1$. We can then use a standard coupling argument so that $N_v(\mu) \geq 1$ if and only if v is not well behaved. The intuition is that, by replacing each node of Π_0 that is not well behaved by a Poisson number of nodes, we can exploit the thinning property of Poisson random variables to show that, as the nodes move, they are stochastically dominated by a Poisson point process.

For each $w \in \Psi_0(\mu)$, let $\xi_w(t)$ be the position of w at time t according to the density function g_t . Define $\Psi_t(\mu)$ to be the point process obtained by

$$\Psi_t(\mu) = \bigcup_{w \in \Psi_0(\mu)} \xi_w(t).$$

The following lemma gives that $\Psi_t(\mu)$ is stochastically dominated by a Poisson point process.

Lemma 2.3. *Let $e^{-\mu}$ be the probability that a node of Π_0 is well behaved. For t sufficiently large, there exists a universal constant $c > 0$ such that, if $\tilde{\Psi}$ is a Poisson point process with intensity $c\sqrt{\delta}$, then it is possible to couple $\tilde{\Psi}$ with $\Psi_t(\mu)$ so that $\Psi_t(\mu) \subseteq \tilde{\Psi}$.*

Proof. Since the nodes of $\Phi_0(\mu)$ move independently of one another, we can apply the thinning property of Poisson random variables to obtain that $\Psi_t(\mu)$ is a Poisson point process. Let $\Lambda(x)$ be the intensity of $\Psi_t(\mu)$ at $x \in \mathbb{R}^2$. By symmetry of Brownian motion and the symmetry in the motion of well behaved nodes, we have that

$$\Lambda(x) = \sum_{v \in \Pi_0} \mu g_t(v, x) = \sum_{v \in \Pi_0} \mu g_t(x, v). \quad (6)$$

Recall that, for any $z \in \mathbb{R}^2$ and $\ell > 0$, we define $z + S_\ell$ as the translation of the square $[0, \ell]^2$ so that its center is at z . Define the square R_1 as $x + S_{5\delta\sqrt{t}}$, the annulus R_2 as $(x + S_{5C\delta\sqrt{t}}) \setminus R_1$ and the region R_3 as $\mathbb{R}^2 \setminus (R_1 \cup R_2)$. We split the sum in (6) into three parts by considering the set of points $P_1 = \Pi_0 \cap R_1$, $P_2 = \Pi_0 \cap R_2$ and $P_3 = \Pi_0 \cap R_3$.

We start with P_2 . We can partition each hexagon of $\mathcal{H}$ into smaller hexagons of side length $\sqrt{3}/3$ such that each point of Π_0 is contained in exactly one such hexagon. This is possible since the dual lattice¹ of $\mathcal{T}$ is a hexagonal lattice of side length $\sqrt{3}/3$, so the hexagons of side length $\sqrt{3}/3$ mentioned above can be obtained by translating and rotating the dual lattice of $\mathcal{T}$. We denote by $\mathcal{H}'$ the set of hexagons of side length $\sqrt{3}/3$ obtained in this way.

For each $z \in \mathbb{R}^2$, let H_z be the hexagon that contains z in $\mathcal{H}'$. Each H_z has side length $\sqrt{3}/3$ and area $\sqrt{3}/2$. Thus, for any point $z \in R_2$, we have that $H_z \subset x + S_{5C\delta\sqrt{t}+4\sqrt{3}/3}$. Let $R'_2 = (x + S_{5C\delta\sqrt{t}+4\sqrt{3}/3}) \setminus R_1$, which gives that

$$\sum_{v \in P_2} \mu g_t(x, v) \leq \frac{2}{\sqrt{3}} \int_{R'_2} \sup_{z' \in H_z} \mu g_t(x, z') dz.$$

Now, note that $\frac{\mu}{\mathbf{P}(v \text{ is not well behaved})} = \frac{\mu}{1-e^{-\mu}} \leq \frac{1}{1-\mu/2} \leq 2$ since $\mu \leq 1$. Then, using the definition of g_t from (5) and the definition of φ_t in (1), we have that

$$\begin{aligned} \sum_{v \in P_2} \mu g_t(x, v) &\leq \frac{4}{\sqrt{3}} \int_{R'_2} \left(\sup_{z' \in H_z} f_t(z' - x) - \varphi_t(q(x), q(z')) \right) dz \\ &\leq \frac{4}{\sqrt{3}} \int_{R'_2} \left(\sup_{z' \in H_z} f_t(z' - x) - \inf_{x' \in Q_{q(x)}, z'' \in Q_{q(z')}} f_t(x' - z'') \right) dz. \end{aligned}$$

Now, by the triangle inequality, we have that $\|z' - x\|_2 \geq \|z - x\|_2 - \|z - z'\|_2 \geq \|z - x\|_2 - 2\sqrt{3}/3$ and $\|x' - z''\|_2 \leq \|z - x\|_2 + \|z - z''\|_2 + \|x - x'\|_2 \leq \|z - x\|_2 + 4\delta\sqrt{t}$, which gives that

$$\begin{aligned} \sum_{v \in P_2} \mu g_t(x, v) &\leq \frac{4}{\sqrt{3}} \int_{R'_2} \frac{1}{2\pi t} \left(\exp \left(-\frac{(\|z - x\|_2 - 2\sqrt{3}/3)^2}{2t} \right) - \exp \left(-\frac{(\|z - x\|_2 + 4\delta\sqrt{t})^2}{2t} \right) \right) dz. \end{aligned}$$

Note that we can write

$$\begin{aligned} &\exp \left(-\frac{(\|z - x\|_2 - 2\sqrt{3}/3)^2}{2t} \right) - \exp \left(-\frac{(\|z - x\|_2 + 4\delta\sqrt{t})^2}{2t} \right) \\ &= \left(\exp \left(\frac{2\sqrt{3}\|z - x\|_2 - 2}{3t} \right) - \exp \left(-\frac{(4\|z - x\|_2\delta\sqrt{t} + 8\delta^2 t)}{t} \right) \right) \exp \left(-\frac{\|z - x\|_2^2}{2t} \right). \end{aligned}$$

Now we use that, for $z \in R'_2$, we have $\|z - x\|_2 \leq \frac{5\sqrt{2}C}{2}\delta\sqrt{t} + 2\sqrt{6}/3$. Then, the first exponential term above is $1 + o(1)$ and, for the second exponential term, we can use the inequality $e^{-x} \geq 1 - x$, which gives, as $t \rightarrow \infty$,

$$\begin{aligned} \sum_{v \in P_2} \mu g_t(x, v) &\leq \frac{4}{\sqrt{3}} (10\sqrt{2}C\delta^2 + 8\delta^2 + o(1)) \int_{R'_2} \frac{1}{2\pi t} \exp \left(-\frac{\|z - x\|_2^2}{2t} \right) dz \\ &\leq c_1\sqrt{\delta} + o(1), \end{aligned} \tag{7}$$

¹Recall that the dual lattice of $\mathcal{T}$ is the lattice whose points are the faces of $\mathcal{T}$ and two points are adjacent if their corresponding faces in $\mathcal{T}$ have a common edge.

for some universal constant $c_1 > 0$.

For the terms of (6) where $v \in P_3$ we have that $g_t(x, v) = \frac{f_t(x, v)}{\mathbf{P}(v \text{ is not well behaved})}$. Then, let $R'_3 = \mathbb{R}^2 \setminus (x + S_{5C\delta\sqrt{t}-4\sqrt{3}/3})$ so that, for each $z \in R_3$, we have $H_z \subset R'_3$, which allows us to write

$$\sum_{v \in P_3} \mu g_t(x, v) \leq \frac{4}{\sqrt{3}} \int_{R'_3} \sup_{z' \in H_z} f_t(x, z') dz \leq \frac{4}{\sqrt{3}} \int_{R'_3} \frac{1}{2\pi t} \exp\left(-\frac{(\|z - x\|_2 - 2\sqrt{3}/3)^2}{2t}\right) dz.$$

Now, letting $w = z - x$ and writing $w = (w_1, w_2)$ we have that

$$(\|w\|_2 - 2\sqrt{3}/3)^2 \geq (|w_1| - 2\sqrt{3}/3)^2 + (|w_2| - 2\sqrt{3}/3)^2 - 4/3,$$

which can be used to get the bound

$$\sum_{v \in P_3} \mu g_t(x, v) \leq \frac{4}{\sqrt{3}} \exp\left(\frac{2}{3t}\right) \left(2 \int_{\frac{5C\delta\sqrt{t}}{2} - \frac{4\sqrt{3}}{3}}^{\infty} \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{w_1^2}{2t}\right) dw_1\right)^2 \leq \frac{c_2}{C\delta} + o(1), \quad (8)$$

for some universal constant $c_2 > 0$. Finally, for the terms in (6) with $v \in P_1$, we use that $\mu g_t(v, x) \leq 2f_t(v, x) \leq \frac{1}{\pi t}$ for all v, x which gives that

$$\sum_{v \in P_1} \mu g_t(x, v) \leq \frac{1}{\pi t} \frac{2}{\sqrt{3}} \left(5\delta\sqrt{t} + 4\frac{\sqrt{3}}{3}\right)^2 \leq c_3\delta^2 + o(1), \quad (9)$$

for some universal constant $c_3 > 0$ and where $\frac{2}{\sqrt{3}} \left(5\delta\sqrt{t} + 4\frac{\sqrt{3}}{3}\right)^2$ is an upper bound for the number of points in P_1 .

Plugging (7), (8) and (9) into (6) yields

$$\Lambda(x) = \sum_{v \in \Pi_0} \mu g_t(x, v) \leq c_1\sqrt{\delta} + \frac{c_2}{C\delta} + c_3\delta^2 + o(1).$$

□

We now proceed to the proof of Theorem 1.3.

Proof of Theorem 1.3. We start by giving a high-level overview of the proof. First, we assume that all nodes of Π_0 are well behaved. Then we consider a hexagon Q_i of $\mathcal{H}$ and count the number of such well behaved nodes that are inside Q_i at time t . Note that, by the definition of well behaved nodes, given that a node is in Q_i at time t , then its location is uniformly random in Q_i . Therefore, in order to show that they are stochastically dominated by a Poisson point process, it suffices to show that there are at most as many nodes of Π_0 in Q_i at time t as nodes of the Poisson point process. This will happen with a probability that can be made arbitrarily large by setting t large enough. We then use the fact that, since nodes are considered well behaved, a node can only be in Q_i at time t if that node was inside a hexagon of J_i at time 0. Therefore, if we consider a hexagon Q_j such that $J_i \cap J_j = \emptyset$, we have that the well-behaved nodes that are able to be in Q_i at time t cannot end up in Q_j . Hence, the event that the well-behaved nodes in Q_i are stochastically dominated by a Poisson point process is independent of the event that the nodes in Q_j are stochastically dominated by a Poisson point process. This bounded dependency is enough to complete the analysis of well behaved nodes. On the other hand, to handle nodes that are not well behaved, we add a discrete Poisson point process at each node of Π_0 so that the probability that we add at least one node at a given $v \in \Pi_0$ is exactly the same as the probability that v is

not well behaved. Thus, this discrete Poisson point process contains the set of nodes that are not well behaved. We then use Lemma 2.3 to conclude that the nodes that are not well behaved are stochastically dominated by a Poisson point process, which concludes the proof.

We now proceed to the rigorous argument. For each $v \in \Pi_0$, let $\xi'_v(t)$ be the position of v at time t given that v is well behaved, and let

$$\Pi'_t = \bigcup_{v \in \Pi_0} \xi'_v(t).$$

Note that, since $e^{-\mu}$ is the probability that a node is well behaved and $\Psi_0(\mu)$ is the point process obtained by adding a random number of nodes to the points of Π_0 according to a Poisson random variable with mean μ , then there exists a coupling so that

$$\Pi_t \subseteq \Pi'_t \cup \Psi_t(\mu).$$

Lemma 2.3 establishes that $\Psi_t(\mu)$ is stochastically dominated by a Poisson point process with intensity $c_1\sqrt{\delta}$ for some universal constant $c_1 > 0$. It remains to show that Π'_t is also stochastically dominated by a Poisson point process. Unfortunately, this is not true in the whole of $\mathbb{R}^2$ as shown in Lemma 2.1. We will then consider the tessellation given by $\mathcal{H}$ and show that, for each hexagon Q_i of the tessellation with $X_i = 1$, where the X_i are defined in the paragraph preceding Theorem 1.3, Π'_t is stochastically dominated by a Poisson point process $\tilde{\Pi}$ of intensity $(1 + \sqrt{\delta})2/\sqrt{3}$.

In order to see this, for each $i \in I$, we define a binary random variable Y_i , which is 1 if $\tilde{\Pi}$ has more nodes in Q_i than Π'_t . Then, since each node of Π'_t is well behaved, whenever $Y_i = 1$, we can couple $\tilde{\Pi}$ with Π'_t such that $\tilde{\Pi} \supseteq \Pi'_t$ in Q_i . First we derive a bound for the number of nodes of Π'_t inside Q_i . For each $v \in \Pi_0$, let Z_v be the indicator random variable for $\xi'_v(t) \in Q_i$. Then, since the probability that $\xi'_v(t) \in Q_i$ is proportional to $\varphi_t(q(v), i)$, the expected number of nodes of Π'_t in Q_i is

$$\sum_{v \in \Pi_0 \cap (\cup_{j \in J_i} Q_j)} \mathbf{E}[Z_v] = \sum_{v \in \Pi_0 \cap (\cup_{j \in J_i} Q_j)} \frac{\varphi_t(q(v), i)}{M} = \sum_{v \in \Pi_0 \cap (\cup_{j \in J_i} Q_j)} \frac{\varphi_t(i, q(v))}{M} = 3\delta^2 t,$$

where M is a normalizing constant so that $\sum_j \varphi_t(i, j) = M$ for all i . The last step follows since $\sum_{v \in \Pi_0 \cap (\cup_{j \in J_i} Q_j)} \varphi_t(i, q(v)) = 3\delta^2 t \sum_{j \in J_i} \varphi_t(i, j) = 3\delta^2 t M$. A simpler way to establish the equation above is by using stationarity and noting that $3\delta^2 t$ is the number of points of Π_0 in Q_i . Since the random variables Z_v are mutually independent, we can apply a Chernoff bound for Binomial random variables (cf. Lemma A.2) to get

$$\mathbf{P} \left(\sum_{v \in \Pi_0 \cap (\cup_{j \in J_i} Q_j)} Z_v \geq (1 + \sqrt{\delta}/2)3\delta^2 t \right) \leq \exp \left(-\frac{2(\sqrt{\delta}/2)^2(3\delta^2 t)^2}{3\delta^2 t |J_i|} \right) \leq \exp \left(-\frac{9\delta^3 t}{8C^2} \right),$$

where the last step follows from Lemma 2.2. Using a standard Chernoff bound for Poisson random variables (cf. Lemma A.1) we have

$$\mathbf{P} \left(\tilde{\Pi} \text{ has less than } (1 + \sqrt{\delta}/2)3\delta^2 t \text{ nodes in } Q_i \right) \leq \exp \left(-\frac{\delta(3\delta^2 t)}{2(1 + \sqrt{\delta})} \right).$$

Therefore, we obtain a constant c_2 such that

$$\mathbf{P}(Y_i = 1) \geq 1 - \exp \left(-\frac{c_2 \delta^3 t}{C^2} \right). \quad (10)$$

The random variables Y are not mutually independent. However, note that Y_i depends only on the random variables $Y_{i'}$ for which $J_{i'} \cap J_i \neq \emptyset$. This is because, for any $i \in I$, only the nodes that are

inside hexagons Q_j with $j \in J_i$ can contribute to Y_i . Therefore, using Lemma 2.2, we have that Y_i depends on at most $(\frac{4}{3}C^2)^2$ other random variables Y . By having t large enough, we can make the bound in (10) be arbitrarily close to 1. This allows us to apply a result of Liggett, Schonmann and Stacey [7, Theorem 1.3], which gives that the random field $(Y_i)_{i \in I}$ stochastically dominates a field $(Y'_i)_{i \in I}$ of independent Bernoulli random variables satisfying

$$\mathbf{P}(Y'_i = 1) \geq 1 - \exp\left(-\frac{c_3 \delta^3 t}{C^6}\right),$$

for some positive constant c_3 . So, with t sufficiently large, we can assure that $\mathbf{P}(Y'_i = 1)$ is larger than p in the statement of Theorem 1.3. Then, we have that, whenever $Y'_i = 1$, the Poisson point process $\tilde{\Pi} \cup \Psi_t(\mu)$ stochastically dominates Π_t inside Q_i . Since $\tilde{\Pi}$ and $\Psi_t(\mu)$ are independent Poisson point processes, we have that their union is also a Poisson point process of intensity no larger than

$$\frac{2}{\sqrt{3}} + \frac{2}{\sqrt{3}}\sqrt{\delta} + c_1\sqrt{\delta}, \quad (11)$$

which completes the proof of Theorem 1.3. $\square$

3 Large time

In this section we give the proofs of Theorems 1.1 and 1.2. Here we will use some steps of the proof of Theorem 1.3 without repeating the details. For this reason, we suggest the reader to read the proof of Theorem 1.3 before embarking in the proofs in this section.

Proof of Theorem 1.1. For each $i \in I$, let N_i be the set of hexagons Q_j such that Q_i and Q_j intersect. Now, define a binary random variable $\tilde{Y}_i$ to be 1 if $Y_j = 1$ for all $j \in N_i$ and the largest component of $R((\cup_{j \in N_i} Q_j) \cap (\tilde{\Pi} \cup \Psi_t(\mu)))$ has diameter smaller than $\delta\sqrt{t}/10$. (Recall the definition of Y_i from the paragraph preceding (10).) If $\lambda_c > \frac{2}{\sqrt{3}}$, we can set δ small enough so that the intensity of $\tilde{\Pi} \cup \Psi_t(\mu)$, which is bounded above by (11), is smaller than λ_c . Then, using (10) and a result of Penrose and Pisztor [11, Theorem 2], we have that, for any given i and sufficiently large t ,

$$\mathbf{P}(\tilde{Y}_i = 1) \geq 1 - 6 \exp\left(-\frac{c_2 \delta^3 t}{C^2}\right) - \exp\left(-c_4 \delta \sqrt{t}\right), \quad (12)$$

for some universal constant $c_4 > 0$. Also, the $\tilde{Y}_i$ depends only on the random variables $Y_{i'}$ for which $J_{i'} \cap J_j \neq \emptyset$ for all $j \in N_i$. Hence, $\tilde{Y}_i$ depends on no more than $7(\frac{4}{3}C^2)^2$ other $\tilde{Y}_j$ since $|N_i| = 7$. Thus, as in the proof of Theorem 1.3, we can apply the result of Liggett, Schonmann and Stacey [7, Theorem 1.3] to conclude that $(\tilde{Y}_i)_i$ stochastically dominates a random field of independent Bernoulli random variables with mean $\tilde{p}$, which can be made arbitrarily close to 1 by having t large enough. As a consequence, we have that, almost surely, the set of hexagons with $\tilde{Y}_i = 0$ has only finite components. With this, we can conclude that $R(\Pi_t)$ has no infinite component almost surely since the coupling between Π_t , $(\tilde{Y}_i)_i$, $\tilde{\Pi}$ and $\Psi_t(\mu)$ gives that a connected component of $R(\Pi_t)$ can only intersect two non-adjacent edges of an hexagon $i \in I$ if $\tilde{Y}_i = 0$. But we showed that any set of intersecting hexagons with $\tilde{Y}_i = 0$ is finite and, therefore, must be surrounded by hexagons j with $\tilde{Y}_j = 1$, which are not crossed by $R(\Pi_t)$. Hence, all components of $R(\Pi_t)$ are finite. $\square$

Proof of Theorem 1.2. In this proof we will use some steps from the proof of Theorem 1.1. Let r_c^λ be the critical radius for percolation of the Boolean model with intensity λ . Therefore, if t is

large enough and we add balls of radius $r < r_c^{\lambda_0}$ centered at the points of Π_t , where λ_0 is the intensity of $\tilde{\Pi} \cup \Psi_t(\mu)$, which is bounded above by (11), we have that (12) holds and the union of the balls do not have an infinite component. Now, we have that the Boolean model with intensity λ and radius r is equivalent (up to scaling) to the Boolean model with intensity $\bar{\lambda}$ and $\bar{r}$ provided $\lambda r^2 = \bar{\lambda} \bar{r}^2$. Therefore, for any $\epsilon > 0$, we have that

$$r_c^{\lambda+\epsilon} = r_c^\lambda \sqrt{\frac{\lambda}{\lambda+\epsilon}}. \quad (13)$$

Using this we obtain that, for any $\delta > 0$,

$$\liminf_{t \rightarrow \infty} r_c(t) \geq r_c^{\lambda_0} = r_c^{2/\sqrt{3}} \sqrt{\frac{2/\sqrt{3}}{2\sqrt{3} + c\sqrt{\delta}}},$$

where $c > 0$ is a universal constant. Since δ can be arbitrarily close to 0, we obtain

$$\liminf_{t \rightarrow \infty} r_c(t) \geq r_c^{2/\sqrt{3}}. \quad (14)$$

Now, to obtain an upper bound for $r_c(t)$, we use a result of Sinclair and Stauffer [13, Proposition 4.1], which can be stated as follows.

Proposition 3.1 ([13, Proposition 4.1]). *Consider a square S of side length K tessellated into subsquares of side length ℓ , and assume that each subsquare contains at least $\beta\ell^2$ nodes at time 0. Denote the nodes by Ξ_0 and let Ξ_Δ be the point process obtained after the nodes of Ξ_0 have moved as independent Brownian motions for time Δ . Then, there are positive constants c_1, c_2, c_3 such that, for any $\epsilon > 0$, if $\Delta \geq c_1\ell^2/\epsilon^2$, we can couple Ξ_0 with an independent Poisson point process of intensity $(1-\epsilon)\beta$ so that, inside a square S' of side length $K' \leq K - c_2\sqrt{\Delta \log \epsilon^{-1}}$ with the same center as S , Ξ_Δ contains the nodes of the Poisson point process with probability at least $1 - e^{-c_3\epsilon^2\beta\ell^2}$.*

Now we show how to apply this result to our setting. Let $i \in I$ be fixed. Take S to be the union of the hexagons in J_i ; clearly S is not a square but that is not important. Instead of tessellating S into squares, we tessellate S into hexagons of side length $\sqrt{3}/3$. Each such hexagon contains one node of $\Xi_0 = \Pi_0 \cap (\cup_{j \in J_i} Q_j)$, which gives $\beta = 2/\sqrt{3}$, the density of nodes of Ξ_0 . The main step in adapting the proof to our setting is to note that ℓ can represent the diameter of the cells of the tessellation, which in our case gives $\ell = 2\sqrt{3}/3$, and $K - K'$ is the minimum distance between S' and a point outside S . Since we take $S' = Q_i$ we have that $K - K' \geq C\delta\sqrt{t} - 2\delta\sqrt{t}$, where $2\delta\sqrt{t}$ is the diameter of Q_i . Now setting $\epsilon = \sqrt{\delta}$ and $\Delta = t$, we obtain a t_0 so that, for all $t \geq t_0$, the conditions on Δ and $K - K'$ in Proposition 3.1 are satisfied. Therefore, with probability at least $1 - e^{-c_3\delta^3 t}$, for some positive constant c_3 , the nodes of Π_t that are inside Q_i at time t and were inside $\cup_{j \in J_i} Q_j$ at time 0 stochastically dominate a Poisson point process Φ of intensity $(1 - \sqrt{\delta})\frac{2}{\sqrt{3}}$. When this happens, we let $Y_i = 1$, where Y_i here is analogous to the one in (10).

Then we proceed similarly as in the proof of Theorem 1.1. Define $\tilde{Y}_i$ to be 1 if $Y_j = 1$ for all $j \in N_i$ (recall that N_i is the set of hexagons that intersect Q_i) and the largest component of the region $R(\Phi) \cap (\cup_{j \in N_i} Q_j)$, which we denote by X , is such that $\cup_{j \in N_i} Q_j \setminus X$ contains only components of diameter smaller than $\delta\sqrt{t}/10$. This means that if there exists a path $\tilde{Y}_1, \tilde{Y}_2, \dots$ such that, for all $j = 1, 2, \dots$, we have $\tilde{Y}_j = 1$ and Q_j and Q_{j+1} intersect, then the region $R(\Phi) \cap (Q_1 \cup Q_2 \cup \dots)$ has a connected component that intersects each hexagon Q_j in the path. Then, for t sufficiently large, we can apply [11, Theorem 2] and [7, Theorem 1.3] as before to show that the hexagons with $\tilde{Y}_i = 1$ percolate and, consequently, adding balls of radius $r > r_c^{2/\sqrt{3}-\sqrt{\delta}}$ centered at the nodes of Φ produces

an infinite component. By the scaling argument in (13) we have that $r_c^{2/\sqrt{3}-\sqrt{\delta}} = r_c^{2/\sqrt{3}} \sqrt{\frac{\lambda}{\lambda-\sqrt{\delta}}}$, which finally yields

$$\limsup_{t \rightarrow \infty} r_c(t) \leq r_c^{2/\sqrt{3}-\sqrt{\delta}} = r_c^{2/\sqrt{3}} \sqrt{\frac{\lambda}{\lambda-\sqrt{\delta}}}.$$

Since δ can be arbitrarily close to 0, we obtain $\limsup_{t \rightarrow \infty} r_c(t) \leq r_c^{2/\sqrt{3}}$, which together with (14) concludes the proof of Theorem 1.2. $\square$

4 Short time

Now we turn our attention to the case when t is sufficiently small. We establish that, given a Monte Carlo estimate, $R(\Pi_t)$ contains an infinite component almost surely for sufficiently small t .

Consider a tessellation of $\mathbb{R}^2$ into regular hexagons of side length 50. We will denote this tessellation by $\mathcal{H}_{50}$. Instead of considering the usual tessellation, where each hexagon is obtained by the union of some triangles of $\mathcal{T}$, we will shift the hexagonal tessellation (see the illustration in Figure 1) so that no node of Π_0 is on an edge or vertex of $\mathcal{H}_{50}$, and the edges of $\mathcal{H}_{50}$ intersect as many of the balls centered at Π_0 as possible. More formally, since a transitive lattice can be specified by a single edge, we define $\mathcal{T}$ as the triangular lattice containing an edge between the points $(0,0)$ and $(1,0)$, and for any $\ell > 0$, we let $\mathcal{H}_\ell$ be the hexagonal lattice containing an edge between $(1/2, -\sqrt{3}/4)$ and $(\ell + 1/2, -\sqrt{3}/4)$.

Let H_1 and H_2 be two hexagons of $\mathcal{H}_{50}$ that have one edge in common, and denote this edge by e . Starting from e , denote the other edges of H_1 in clockwise direction by e_1, e_2, e_3, e_4, e_5 ; thus e_3 is the edge of H_1 opposite to e . Similarly, denote the other edges of H_2 in clockwise direction by $e'_1, e'_2, e'_3, e'_4, e'_5$ (refer to Figure 1). Given any three sets $X_1, X_2, X_3 \subset \mathbb{R}^2$ and any $t > 0$, we say that $R(\Pi_t)$ has a path from X_1 to X_2 inside X_3 if there exists a sequence of nodes $u_1, u_2, \dots, u_k$ of Π_t , all of which inside X_3 , such that $B(u_1, 1/2)$ intersects X_1 , $B(u_k, 1/2)$ intersects X_2 , and for each $i \geq 1$, the distance between u_i and u_{i+1} is at most 1. With this, we say that $R(\Pi_t)$ crosses H_1 and H_2 if the following three conditions hold:

1. $R(\Pi_t)$ has a path from e_3 to e'_3 inside $H_1 \cup H_2$.
2. $R(\Pi_t)$ has a path from $e_1 \cup e_2$ to $e_4 \cup e_5$ inside $H_1 \cup H_2$.
3. $R(\Pi_t)$ has a path from $e'_1 \cup e'_2$ to $e'_4 \cup e'_5$ inside $H_1 \cup H_2$.

We denote by A_t the event that $R(\Pi_t)$ crosses H_1 and H_2 with a path that also crosses H_1 and H_2 at time 0; this last condition is used to obtain a type of monotonicity later. Then we have the following result.

Theorem 4.1. *Suppose that there exists an $\epsilon_0 > 0$ such that $\mathbf{P}(A_{\epsilon_0}) > 0.8639$. Then, for all $\epsilon \in [0, \epsilon_0]$, it holds that $R(\Pi_\epsilon)$ contains an infinite connected component almost surely.*

Note that, for any fixed t , verifying the condition $\mathbf{P}(A_t) > 0.8639$ resorts to solving a finite, but high dimensional integral describing the crossing probability. We were able to check the validity of this condition for $t = 0.01$ via a Monte Carlo analysis² with confidence 99.99%.

We start the proof with the lemma below, which establishes a type of monotonicity that will be useful later. To state the lemma, let V be a set of points in $\mathbb{R}^d$ and define $E(V)$ as the set of pairs of points of V whose distance is at most 1. Note that the pair $(V, E(V))$ induces a graph over V .

²To obtain this Monte Carlo estimate we employed the Mersenne Twister pseudorandom number generator by Matsumoto and Nishimura [8] with period $2^{19937} - 1$ and improved initialization scheme from January 26th, 2002.




Figure 1: The hexagonal tessellation $\mathcal{H}_4$ of $\mathbb{R}^2$ with hexagons of side length 4, and the illustration of a path of intersecting balls crossing H_1 and H_2 .

Lemma 4.2. *Let $V \subseteq \Pi_0$ such that the graph $(V, E(V))$ is connected. Let V_s be obtained by letting the nodes of V move for time $s \geq 0$ according to independent Brownian motions. Then, we have that $\mathbf{P}(E(V) \subseteq E(V_s))$ is non-increasing with s .*

Proof. This follows by Brownian scale. Consider $s' > s$, and let $V_{s'}$ be obtained by letting the nodes of V move according to independent Brownian motions for time s' . Now we create a coupling between $E(V_{s'})$ and $E(V_s)$ so that, if $E(V) \subseteq E(V_{s'})$, then $E(V) \subseteq E(V_s)$.

For each $u \in V$, let $\zeta_u(s)$ and $\zeta'_u(s')$ be the Brownian motions for the motion of u in V_s and $V_{s'}$, respectively. Then,

$$\mathbf{P}(E(V) \subseteq E(V_s)) = \mathbf{P}\left(\bigcap_{(u,v) \in E(V)} \left\{\|u + \zeta_u(s) - v - \zeta_v(s)\|_2 \leq 1\right\}\right).$$

Now, by Brownian scale, we can couple $\zeta_u(s)$ and $\zeta'_u(s')$ via $\zeta_u(s) = \sqrt{s/s'}\zeta'_u(s')$. Using this, we write the right-hand side above as

$$\mathbf{P}\left(\bigcap_{(u,v) \in E(V)} \left\{\|u - v + \sqrt{s/s'}(\zeta'_u(s') - \zeta'_v(s'))\|_2 \leq 1\right\}\right).$$

Now define the vectors $x_1 = u - v$ and $x_2 = \zeta'_u(s') - \zeta'_v(s')$. For any $\delta \in (0, 1)$, it follows by standard geometric arguments that, if $\|x_1\|_2$ and $\|x_1 + x_2\|_2$ are at most 1, then

$$\|x_1 + \delta x_2\|_2 \leq 1,$$

which establishes that

$$\begin{aligned} & \mathbf{P}\left(\bigcap_{(u,v) \in E(V)} \left\{\|u - v + \sqrt{s/s'}(\zeta'_u(s') - \zeta'_v(s'))\|_2 \leq 1\right\}\right) \\ & \geq \mathbf{P}\left(\bigcap_{(u,v) \in E(V)} \left\{\|(u - v) + (\zeta'_u(s') - \zeta'_v(s'))\|_2 \leq 1\right\}\right) = \mathbf{P}(E(V) \subseteq E(V_{s'})), \end{aligned}$$

which completes the proof. $\square$

Now we proceed to the proof of Theorem 4.1.

Proof of Theorem 4.1. Note that, due to Lemma 4.2, proving Theorem 4.1 reduces to showing that, if there exists an $\epsilon > 0$ such that $\mathbf{P}(A_\epsilon) > 0.8639$, then $R(\Pi_\epsilon)$ contains an infinite connected component almost surely. We henceforth fix a value of ϵ and assume that $\mathbf{P}(A_\epsilon) > 0.8639$.

We will use a renormalization argument. Consider the hexagons $\mathcal{H}_{50}$ described in the beginning of this section. Now, define the graph $L = (U, F)$ such that U is the set of points given by the centers of the hexagons and F is the set of edges between every pair of points $i, j \in U$ for which the hexagons with centers at i and j share an edge. Note that L consists of a scaling of the triangular lattice.

We now define a collection of random variables X_i for each edge $i \in F$. In order to explain the process defining X_i , let H_1 and H_2 be the hexagons whose centers are the endpoints of i . We then define $X_i = 1$ if and only if $R(\Pi_\epsilon)$ crosses H_1 and H_2 with a path of balls that also crosses H_1 and H_2 at time 0. (The definition of crossings is given right before the statement of Theorem 4.1.) Let j be an edge such that i and j are disjoint, and let H_3 and H_4 denote the hexagons centered at the endpoints of j . Clearly, X_i and X_j are independent since the set of balls crossing H_1 and H_2 at time 0 does not intersect the set of balls crossing H_3 and H_4 at time 0. Thus, the collection $(X_i)_i$ is a so-called 1-dependent bond percolation process, with $\mathbf{P}(X_i = 1) = \mathbf{P}(A_\epsilon) > 0.8639$. Then, we can use a result of Balister, Bollobás and Walters [2, Theorem 2], which gives that any 1-dependent bond percolation process on the square lattice with marginal probability larger than 0.8639 percolates almost surely. Since the triangular lattice contains the square lattice, we obtain that, almost surely, there exists an infinite path of consecutive edges of F with $X_i = 1$ for all i in the path.

To conclude the proof, note that, for two non-disjoint edges i and j with $X_i, X_j = 1$, we have that the crossings of the hexagons whose centers are located at the endpoints of i and j intersect. Then, the infinite path of X_i with $X_i = 1$ for all i in the path contains an infinite path inside $R(\Pi_\epsilon)$, which concludes the proof of Theorem 4.1. $\square$

5 Extensions and Open Problems

In the remaining of this section we discuss extensions and open problems regarding other circle packings (Section 5.1), balls moving over graphs (Section 5.2) and critical radius for non-mobile point processes (Section 5.3)

5.1 Other circle packings

Let Π_0^s be the point process given by the vertices of the square lattice with side length 1, and let Π_t^s be the point process obtained by letting the nodes of Π_0^s move for time t according to independent Brownian motions. Note that, for any $\epsilon > 0$, if we look at two balls of radius $1/2$ centered at two adjacent nodes of Π_0^s , then at time ϵ , the probability that these two balls intersect is strictly smaller than $1/2$, which is the critical probability for edge percolation on the square lattice [5]. This motivates our next conjecture.

Conjecture 5.1. *For any $\epsilon > 0$, it holds that, almost surely, all components of $R(\Pi_\epsilon^s)$ are finite.*

Now we consider the question of whether percolation is a *monotone* property. We say that a point process Π_0 is transitive if, for every two nodes $v, v' \in \Pi_0$, there exists an isometry $f: \Pi_0 \rightarrow \Pi_0$ such that $f(v) = v'$. The open problem below concerns the question of whether transitivity is enough to obtain monotonicity in the percolation properties of balls moving as Brownian motion.




Figure 2: An example of a non-transitive configuration of balls that is not monotone. Each solid ball represents the superposition of 14 balls and white balls represent single balls.

Question 5.2. Let Π_0 be a transitive point process so that $R(\Pi_0)$ is a connected set. Let Π_t be obtained from Π_0 by letting the nodes move as independent Brownian motions for time t . Then, if for some time t_0 we have that $R(\Pi_{t_0})$ has an infinite component almost surely, then, is it true that, for any $t < t_0$, $R(\Pi_t)$ also has an infinite component almost surely? Similarly, if for some t_1 we have that $R(\Pi_{t_1})$ contains only finite components almost surely, then, does it hold that, for any $t > t_1$, $R(\Pi_t)$ also contains only finite components almost surely?

Remark 5.3. We note that Question 5.2 above is false if we drop the condition that Π_0 is transitive. For example, consider a tessellation of $\mathbb{R}^2$ into squares of side length 6 and, in each square of the tessellation, consider the configuration of balls illustrated in Figure 2, where each ball has radius $1/2$, solid balls represent the superposition of 14 balls and white balls represent a single ball. It is easy to see that, at a sufficiently small time ϵ , the union of the balls will not contain an infinite component almost surely. However, the density of balls is equal to $\frac{9 \times 14 + 18}{36} = 4$ and, as the balls move for a sufficiently large amount of time, their position will approach a Poisson point process which is known to percolate.

5.2 Motion over graphs

We now consider the case when the motion of the nodes is more restricted. First, let Π_0 be the point process given by the integer points of $\mathbb{R}$. For any node $u \in \Pi_0$, we let $u + \zeta_u(t)$ be its position at time t , where $(\zeta_u(t))_t$ is a one-dimensional Brownian motion. Now, consider a sequence of m distinct nodes $u_1, u_2, \dots, u_m$ such that $B(u_i, 1/2)$ and $B(u_{i+1}, 1/2)$ intersect for all i . We call such a sequence of nodes as a *path*. Let ϵ be a sufficiently small positive constant, and consider only the nodes of Π_0 whose displacement from time 0 to time ϵ is smaller than $1/2$; we denote these nodes as *good* nodes. We claim that

$$\mathbf{P}(u_1, u_2, \dots, u_m \text{ form a path at time } \epsilon \mid u_i \text{ is good for all } i) = \frac{1}{m!}. \quad (15)$$

In order to see this, suppose, without loss of generality, that $u_1 < u_2 < \dots < u_m$. For each node $u \in \Pi_0$, let $\zeta'_u(\epsilon)$ be the displacement of u from time 0 to ϵ given that u is a good node. Then, in order for $B(u_1 + \zeta'_{u_1}(\epsilon), 1/2)$ to intersect $B(u_2 + \zeta'_{u_2}(\epsilon), 1/2)$ we need that $|u_1 + \zeta'_{u_1}(\epsilon) - u_2 - \zeta'_{u_2}(\epsilon)| \leq 1$. Since u_1 and u_2 are good nodes, this condition translates to $u_2 + \zeta'_{u_2}(\epsilon) - u_1 - \zeta'_{u_1}(\epsilon) \leq 1$, which in turn implies that $\zeta'_{u_1}(\epsilon) \geq \zeta'_{u_2}(\epsilon)$. Repeating this argument, we obtain the condition $\zeta'_{u_1}(\epsilon) \geq \zeta'_{u_2}(\epsilon) \geq \zeta'_{u_3}(\epsilon) \geq \dots \geq \zeta'_{u_m}(\epsilon)$. Since the ζ' are independent and identically distributed, we have that $\mathbf{P}(\zeta'_{u_1}(\epsilon) \geq \zeta'_{u_2}(\epsilon) \geq \dots \geq \zeta'_{u_m}(\epsilon)) = 1/m!$, which establishes (15).

We now consider a more general scenario. Let G be an infinite graph that is vertex transitive and has bounded degree. We assume that each edge of G has length 1, which gives a metric over G . Let $\Pi_0(G)$ be the point process given by putting one node at each vertex of G and define $\Pi_t(G)$ as the point process obtained by letting the nodes of $\Pi_0(G)$ move for time t along the edges of G according to independent Brownian motions. Then $R(\Pi_t(G))$ is the union of balls centered at the nodes of Π_t and having radius $1/2$ with respect to the metric induced by G . We note that the probability given in (15) for any fixed path $u_1, u_2, \dots, u_m$ of good nodes to form a path at a time ϵ that is sufficiently small is at most $1/m!$. This motivates our next conjecture.

Conjecture 5.4. *Let G be an infinite graph that is vertex transitive and has bounded degree. Then, for any $t > 0$, the region $R(\Pi_t(G))$ contains only finite components almost surely.*

5.3 Critical radius of point processes

Here we let Π be a point process over $\mathbb{R}^2$ and consider the region $R(\Pi, r)$ as the union of balls of radius r centered at the nodes of Π . In this section, we only consider point processes with unit intensity and let $r_c(\Pi)$ be the smallest r for which $R(\Pi, r)$ contains an infinite component. It is intuitive to believe that point processes that are more organized have smaller critical radius; this is the core of our next conjecture. For more information on zeros of Gaussian analytic functions, we refer to [6].

Conjecture 5.5. *Let Π_L be any transitive point process with intensity 1 (as defined before Question 5.2). Let Π_{GAF} be a point process given by the zeros of a Gaussian analytic function with intensity 1 and Π_P be a Poisson point process with intensity 1. Then,*

$$r_c(\Pi_L) < r_c(\Pi_{\text{GAF}}) < r_c(\Pi_P).$$

Finally, consider a Poisson point process Π with intensity 1 over $\mathbb{R}^d$ and let r_c be the critical radius for percolation of balls centered at the nodes of Π . Our last open problem concerns small perturbations of the critical radius.

Question 5.6. *Let $\epsilon > 0$ and, for each node $v \in \Pi$, let X_v be a uniform random variable over $[-\epsilon, \epsilon]$. For each node $v \in \Pi$, add a ball of radius $r_c + X_v$ centered at v . Will the union of the balls contain an infinite component almost surely?*

References

- [1] N. Alon and J.H. Spencer. *The probabilistic method*. John Wiley & Sons, 3rd edition, 2008.
- [2] P. Balister, B. Bollobás, and M. Walters. Continuum percolation with steps in the square or the disc. *Random structures and algorithms*, 26:392–403, 2005.
- [3] B. Bollobás and O. Riordan. *Percolation*. Cambridge University Press, 2006.

- [4] J.H. Conway and N.J.A. Sloane. *Sphere packings, lattices and groups*. Springer-Verlag, 3rd edition, 1999.
- [5] G. Grimmett. *Percolation*. Springer-Verlag, 2nd edition, 1999.
- [6] J.B. Hough, M. Krishnapur, Y. Peres, and B. Virág. *Zeros of Gaussian Analytic functions and determinantal point processes*. American Mathematical Society, 2009.
- [7] T.M. Liggett, R.H. Schonmann, and A.M. Stacey. Domination by product measures. *The Annals of Probability*, 25:71–95, 1997.
- [8] M. Matsumoto and T. Nishimura. Mersenne twister: A 623-dimensionally equidistributed uniform pseudo-random number generator. *ACM Transactions on Modeling and Computer Simulation*, 8:3–30, 1998.
- [9] R. Meester and R. Roy. *Continuum Percolation*. Cambridge University Press, 1996.
- [10] P. Mörters and Y. Peres. *Brownian Motion*. Cambridge University Press, 2010.
- [11] M. Penrose and A. Pisztor. Large deviations for discrete and continuous percolation. *Advances in Applied Probability*, 28:29–52, 1996.
- [12] Y. Peres, A. Sinclair, P. Sousi, and A. Stauffer. Mobile geometric graphs: detection, coverage and percolation. In *Proceedings of the 22st ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 412–428, 2011.
- [13] A. Sinclair and A. Stauffer. Mobile geometric graphs, and detection and communication problems in mobile wireless networks, 2010. arXiv:1005.1117v2.
- [14] A. Stauffer. Adversarial detection and space-time percolation in mobile geometric graphs, 2011. arXiv:1108.6322v1.

A Standard large deviation results

We use the following standard Chernoff bounds during our proofs.

Lemma A.1 (Chernoff bound for Poisson). *Let P be a Poisson random variable with mean λ . Then, for any $0 < \epsilon < 1$,*

$$\mathbf{P}(P \geq (1 + \epsilon)\lambda) \leq \exp\left(-\frac{\lambda\epsilon^2}{2}(1 - \epsilon/3)\right) \quad \text{and} \quad \mathbf{P}(P \leq (1 - \epsilon)\lambda) \leq \exp\left(-\frac{\lambda\epsilon^2}{2}\right).$$

Lemma A.2 (Chernoff bound for Binomial, see [1, Lemma A.1.4]). *Let $X_1, X_2, \dots, X_n$ be independent Bernoulli random variable such that $\mathbf{E}[X_i] = p_i$. Let $X = \sum_{i=1}^n X_i$. Then, for any $\epsilon > 0$,*

$$\mathbf{P}(X \geq (1 + \epsilon)\mathbf{E}[X]) \leq \exp\left(-\frac{2\epsilon^2(\mathbf{E}[X])^2}{n}\right) \quad \text{and} \quad \mathbf{P}(X \leq (1 - \epsilon)\mathbf{E}[X]) \leq \exp\left(-\frac{\lambda\epsilon^2}{2}\right).$$

Lemma A.3 (Gaussian tail bound [10, Theorem 12.9]). *Let X be a normal random variable with mean 0 and variance σ^2 . Then, for any $R \geq \sigma$ we have that $\mathbf{P}(X \geq R) \leq \frac{\sigma}{\sqrt{2\pi}R} \exp\left(-\frac{R^2}{2\sigma^2}\right)$.*